arXiv:1101.0944v3 [math.CA] 15 Jul 2011

INEQUALITIES FOR CONVEX AND $s-$CONVEX FUNCTIONS ON $\Delta=[a,b]\times[c,d]$

M. EMIN ÖZDEMIR♦, HAVVA KAVURMACI♦,★, AHMET OCAK AKDEMIR♠,
AND MERVE AVCI♦

Abstract. In this paper, two new lemmas are proved and inequalities are established for co-ordinated convex functions and co-ordinated $s-$convex functions.

1. INTRODUCTION

Let $f:I\subseteq\mathbb{R}\to\mathbb{R}$ be a convex function defined on the interval I of real numbers and $a<b$. The following double inequality;

$$f\left(\frac{a+b}{2}\right)\leq\frac{1}{b-a}\int_{a}^{b}f(x)dx\leq\frac{f(a)+f(b)}{2}$$

is well known in the literature as Hadamard's inequality. Both inequalities hold in the reversed direction if f is concave.

In [8], Orlicz defined $s-$convex functions as following:

Definition 1. *A function $f:\mathbb{R}^{+}\to\mathbb{R}$, where $\mathbb{R}^{+}=[0,\infty)$, is said to be $s-$convex in the first sense if*

$$f(\alpha x+\beta y)\leq\alpha^{s}f(x)+\beta^{s}f(y)$$

for all $x,y\in[0,\infty)$, $\alpha,\beta\geq0$ with $\alpha^{s}+\beta^{s}=1$ and for some fixed $s\in(0,1]$. We denote by K_{s}^{1} the class of all $s-$convex functions.

Definition 2. *A function $f:\mathbb{R}^{+}\to\mathbb{R}$, where $\mathbb{R}^{+}=[0,\infty)$, is said to be $s-$convex in the second sense if*

$$f(\alpha x+\beta y)\leq\alpha^{s}f(x)+\beta^{s}f(y)$$

for all $x,y\in[0,\infty)$, $\alpha,\beta\geq0$ with $\alpha+\beta=1$ and for some fixed $s\in(0,1]$. We denote by K_{s}^{2} the class of all $s-$convex functions.

Obviously, one can see that if we choose $s=1$, both definitions reduced to ordinary concept of convexity.

For several results related to above definitions we refer readers to [6], [9], [13].

In [2], Dragomir defined convex functions on the co-ordinates as following:

Date: December 29, 2010.

2000 *Mathematics Subject Classification.* 26D10,26D15.

Key words and phrases. Ostrowski's inequality, co-ordinates, convex functions, $s-$convex functions.

★Corresponding Author.

Definition 3. *Let us consider the bidimensional interval* $\Delta = [a,b] \times [c,d]$ *in* $\mathbb{R}^2$ *with* $a < b$, $c < d$. *A function* $f : \Delta \to \mathbb{R}$ *will be called convex on the co-ordinates if the partial mappings* $f_y : [a,b] \to \mathbb{R}$, $f_y(u) = f(u,y)$ *and* $f_x : [c,d] \to \mathbb{R}$, $f_x(v) = f(x,v)$ *are convex where defined for all* $y \in [c,d]$ *and* $x \in [a,b]$. *Recall that the mapping* $f : \Delta \to \mathbb{R}$ *is convex on* Δ *if the following inequality holds,*

$$f(\lambda x + (1-\lambda)z, \lambda y + (1-\lambda)w) \leq \lambda f(x,y) + (1-\lambda) f(z,w)$$

for all $(x,y),(z,w) \in \Delta$ *and* $\lambda \in [0,1]$.

In [2], Dragomir established the following inequalities of Hadamard's type for co-ordinated convex functions on a rectangle from the plane $\mathbb{R}^2$.

Theorem 1. *Suppose that* $f : \Delta = [a,b] \times [c,d] \to \mathbb{R}$ *is convex on the co-ordinates on* Δ. *Then one has the inequalities;*

$$
\begin{aligned}
(1.1) \qquad & f\left(\frac{a+b}{2}, \frac{c+d}{2}\right) \\
\leq\ & \frac{1}{2}\left[\frac{1}{b-a}\int_a^b f\left(x, \frac{c+d}{2}\right)dx + \frac{1}{d-c}\int_c^d f\left(\frac{a+b}{2}, y\right)dy\right] \\
\leq\ & \frac{1}{(b-a)(d-c)}\int_a^b \int_c^d f(x,y)\,dxdy \\
\leq\ & \frac{1}{4}\left[\frac{1}{(b-a)}\int_a^b f(x,c)dx + \frac{1}{(b-a)}\int_a^b f(x,d)dx \right. \\
& \left. + \frac{1}{(d-c)}\int_c^d f(a,y)dy + \frac{1}{(d-c)}\int_c^d f(b,y)dy\right] \\
\leq\ & \frac{f(a,c)+f(a,d)+f(b,c)+f(b,d)}{4}.
\end{aligned}
$$

The above inequalities are sharp.

Similar results can be found in [5], [7], [10] and [11].

In [10], Alomari and Darus defined co-ordinated $s-$convex functions and proved some inequalities based on this definition.

Definition 4. *Consider the bidimensional interval* $\Delta = [a,b] \times [c,d]$ *in* $[0,\infty)^2$ *with* $a < b$ *and* $c < d$. *The mapping* $f : \Delta \to \mathbb{R}$ *is* $s-$*convex on* Δ *if*

$$f(\lambda x + (1-\lambda)z, \lambda y + (1-\lambda)w) \leq \lambda^s f(x,y) + (1-\lambda)^s f(z,w)$$

holds for all (x,y), $(z,w) \in \Delta$ *with* $\lambda \in [0,1]$ *and for some fixed* $s \in (0,1]$.

In [12], Sarıkaya *et al.* proved some Hadamard's type inequalities for co-ordinated convex functions as followings:

Theorem 2. *Let* $f : \Delta \subset \mathbb{R}^2 \to \mathbb{R}$ *be a partial differentiable mapping on* $\Delta := [a,b] \times [c,d]$ *in* $\mathbb{R}^2$ *with* $a < b$ *and* $c < d$. *If* $\left|\frac{\partial^2 f}{\partial t \partial s}\right|$ *is a convex function on the* *co-ordinates on* Δ*, then one has the inequalities:*

$$\begin{aligned}
&\left|\frac{f(a,c)+f(a,d)+f(b,c)+f(b,d)}{4}\right. \\
&\qquad\left.\frac{1}{(b-a)(d-c)}\int_a^b\int_c^d f(x,y)dxdy - A\right| \\
\leq\ & \frac{(b-a)(d-c)}{16}\left(\frac{\left|\frac{\partial^2 f}{\partial t\partial s}\right|(a,c)+\left|\frac{\partial^2 f}{\partial t\partial s}\right|(a,d)+\left|\frac{\partial^2 f}{\partial t\partial s}\right|(b,c)+\left|\frac{\partial^2 f}{\partial t\partial s}\right|(b,d)}{4}\right)
\end{aligned}\tag{1.2}$$

where

$$A=\frac{1}{2}\left[\frac{1}{(b-a)}\int_a^b\left[f(x,c)+f(x,d)\right]dx+\frac{1}{(d-c)}\int_c^d\left[f(a,y)dy+f(b,y)\right]dy\right].$$

Theorem 3. *Let* $f:\Delta\subset\mathbb{R}^2\to\mathbb{R}$ *be a partial differentiable mapping on* $\Delta:=[a,b]\times[c,d]$ *in* $\mathbb{R}^2$ *with* $a<b$ *and* $c<d$*. If* $\left|\frac{\partial^2 f}{\partial t\partial s}\right|^q$, $q>1$, *is a convex function on the co-ordinates on* Δ*, then one has the inequalities:*

$$\begin{aligned}
&\left|\frac{f(a,c)+f(a,d)+f(b,c)+f(b,d)}{4}\right. \\
&\qquad\left.\frac{1}{(b-a)(d-c)}\int_a^b\int_c^d f(x,y)dxdy - A\right| \\
\leq\ & \frac{(b-a)(d-c)}{4\,(p+1)^{\frac{2}{p}}}\left(\frac{\left|\frac{\partial^2 f}{\partial t\partial s}\right|^q(a,c)+\left|\frac{\partial^2 f}{\partial t\partial s}\right|^q(a,d)+\left|\frac{\partial^2 f}{\partial t\partial s}\right|^q(b,c)+\left|\frac{\partial^2 f}{\partial t\partial s}\right|^q(b,d)}{4}\right)^{\frac{1}{q}}
\end{aligned}\tag{1.3}$$

where

$$A=\frac{1}{2}\left[\frac{1}{(b-a)}\int_a^b\left[f(x,c)+f(x,d)\right]dx+\frac{1}{(d-c)}\int_c^d\left[f(a,y)dy+f(b,y)\right]dy\right]$$

and $\frac{1}{p}+\frac{1}{q}=1$.

Theorem 4. *Let* $f:\Delta\subset\mathbb{R}^2\to\mathbb{R}$ *be a partial differentiable mapping on* $\Delta:=[a,b]\times[c,d]$ *in* $\mathbb{R}^2$ *with* $a<b$ *and* $c<d$*. If* $\left|\frac{\partial^2 f}{\partial t\partial s}\right|^q$, $q\geq 1$, *is a convex function on the co-ordinates on* Δ*, then one has the inequalities:*

$$\begin{aligned}
&\left|\frac{f(a,c)+f(a,d)+f(b,c)+f(b,d)}{4}\right. \\
&\qquad\left.\frac{1}{(b-a)(d-c)}\int_a^b\int_c^d f(x,y)dxdy - A\right| \\
\leq\ & \frac{(b-a)(d-c)}{16}\left(\frac{\left|\frac{\partial^2 f}{\partial t\partial s}\right|^q(a,c)+\left|\frac{\partial^2 f}{\partial t\partial s}\right|^q(a,d)+\left|\frac{\partial^2 f}{\partial t\partial s}\right|^q(b,c)+\left|\frac{\partial^2 f}{\partial t\partial s}\right|^q(b,d)}{4}\right)^{\frac{1}{q}}
\end{aligned}\tag{1.4}$$

where

$$A=\frac{1}{2}\left[\frac{1}{(b-a)}\int_a^b\left[f(x,c)+f(x,d)\right]dx+\frac{1}{(d-c)}\int_c^d\left[f(a,y)dy+f(b,y)\right]dy\right].$$

In [1], Barnett and Dragomir proved an Ostrowski-type inequality for double integrals as following:

Theorem 5. *Let $f:[a,b]\times[c,d]\to\mathbb{R}$ be continuous on $[a,b]\times[c,d]$, $f''_{x,y}=\frac{\partial^2 f}{\partial x\partial y}$ exists on $(a,b)\times(c,d)$ and is bounded, that is*

$$\left\|f''_{x,y}\right\|_\infty=\sup_{(x,y)\in(a,b)\times(c,d)}\left|\frac{\partial^2 f(x,y)}{\partial x\partial y}\right|<\infty,$$

then we have the inequality;

$$\begin{aligned}
&\left|\int_a^b\int_c^d f(s,t)dtds-(b-a)\int_c^d f(x,t)dt\right.\\
&\left.-(d-c)\int_a^b f(s,y)ds-(b-a)(d-c)f(x,y)\right|\\
\leq\ &\left[\frac{(b-a)^2}{4}+\left(x-\frac{a+b}{2}\right)^2\right]\left[\frac{(d-c)^2}{4}+\left(y-\frac{c+d}{2}\right)^2\right]\left\|f''_{x,y}\right\|_\infty
\end{aligned}\tag{1.5}$$

for all $(x,y)\in[a,b]\times[c,d]$.

In [4], Sarıkaya proved an Ostrowski-type inequality for double integrals and gave a corollary as following:

Theorem 6. *Let $f:[a,b]\times[c,d]\to\mathbb{R}$ be an absolutely continuous functions such that the partial derivative of order 2 exist and is bounded, i.e.,*

$$\left\|\frac{\partial^2 f(t,s)}{\partial t\partial s}\right\|_\infty=\sup_{(x,y)\in(a,b)\times(c,d)}\left|\frac{\partial^2 f(t,s)}{\partial t\partial s}\right|<\infty$$

for all $(t,s)\in[a,b]\times[c,d]$. Then we have,

$$\begin{aligned}
&\left|(\beta_1-\alpha_1)(\beta_2-\alpha_2)f\left(\frac{a+b}{2},\frac{c+d}{2}\right)+H(\alpha_1,\alpha_2,\beta_1,\beta_2)+G(\alpha_1,\alpha_2,\beta_1,\beta_2)\right.\\
&-(\beta_2-\alpha_2)\int_a^b f\left(t,\frac{c+d}{2}\right)dt-(\beta_1-\alpha_1)\int_c^d f\left(\frac{a+b}{2},s\right)ds\\
&-\int_a^b\left[(\alpha_2-c)f(t,c)+(d-\beta_2)f(t,d)\right]dt\\
&\left.-\int_c^d\left[(\alpha_1-a)f(a,s)+(b-\beta_1)f(b,s)\right]ds+\int_a^b\int_c^d f(t,s)dsdt\right|\\
\leq\ &\left[\frac{(\alpha_1-a)^2+(b-\beta_1)^2}{2}+\frac{(a+b-2\alpha_1)^2+(a+b-2\beta_1)^2}{8}\right]\\
&\times\left[\frac{(\alpha_2-c)^2+(d-\beta_2)^2}{2}+\frac{(c+d-2\alpha_2)^2+(c+d-2\beta_2)^2}{8}\right]\left\|\frac{\partial^2 f(t,s)}{\partial t\partial s}\right\|_\infty
\end{aligned}\tag{1.6}$$

for all $(\alpha_1,\alpha_2),(\beta_1,\beta_2)\in[a,b]\times[c,d]$ with $\alpha_1<\beta_1$, $\alpha_2<\beta_2$ where

$$\begin{aligned}
&H(\alpha_1,\alpha_2,\beta_1,\beta_2)\\
=\ &(\alpha_1-a)\left[(\alpha_2-c)f(a,c)+(d-\beta_2)f(a,d)\right]\\
&+(b-\beta_1)\left[(\alpha_2-c)f(b,c)+(d-\beta_2)f(b,d)\right]
\end{aligned}$$

and

$$\begin{aligned}
& G(\alpha_1,\alpha_2,\beta_1,\beta_2) \\
= & (\beta_1-\alpha_1)\left[(\alpha_2-c)\,f\left(\frac{a+b}{2},c\right)+(d-\beta_2)\,f\left(\frac{a+b}{2},d\right)\right] \\
& +(\beta_2-\alpha_2)\left[(\alpha_1-a)\,f\left(a,\frac{c+d}{2}\right)+(b-\beta_1)\,f\left(b,\frac{c+d}{2}\right)\right].
\end{aligned}$$

Corollary 1. *Under the assumptions of Theorem 6, we have*

$$\begin{aligned}
& \left|(b-a)(d-c)\,f\left(\frac{a+b}{2},\frac{c+d}{2}\right)+\int_a^b\int_c^d f(t,s)dsdt\right. \\
& \left.-(d-c)\int_a^b f\left(t,\frac{c+d}{2}\right)dt-(b-a)\int_c^d f\left(\frac{a+b}{2},s\right)ds\right| \\
\leq & \frac{1}{16}\left\|\frac{\partial^2 f(t,s)}{\partial t\partial s}\right\|_\infty (b-a)^2(d-c)^2.
\end{aligned} \tag{1.7}$$

In [3], Pachpatte established a new Ostrowski type inequality similar to inequality (1.5) by using elemantery analysis.

The main purpose of this paper is to establish inequalities of Ostrowski-type for co-ordinated convex functions by using Lemma 2 and to establish some new Hadamard's type inequalities for co-ordinated $s-$convex functions.

2. INEQUALITIES FOR CO-ORDINATED CONVEX FUNCTIONS

To prove our main result, we need the following lemma which contains kernels similar to Barnett and Dragomir's kernels in [1], [see the paper [1], proof of Theorem 2.1].

Lemma 1. *Let* $f:\Delta=[a,b]\times[c,d]\to\mathbb{R}$ *be a partial differentiable mapping on* $\Delta=[a,b]\times[c,d]$. *If* $\frac{\partial^2 f}{\partial t\partial s}\in L(\Delta)$, *then the following equality holds:*

$$\begin{aligned}
& f\left(\frac{a+b}{2},\frac{c+d}{2}\right) \\
& -\frac{1}{(d-c)}\int_c^d f\left(\frac{a+b}{2},y\right)dy-\frac{1}{(b-a)}\int_a^b f\left(x,\frac{c+d}{2}\right)dx \\
& +\frac{1}{(b-a)(d-c)}\int_a^b\int_c^d f(x,y)\,dydx \\
= & \frac{1}{(b-a)(d-c)}\int_a^b\int_c^d p(x,t)\,q(y,s)\,\frac{\partial^2 f}{\partial t\partial s}\left(\frac{b-t}{b-a}a+\frac{t-a}{b-a}b,\frac{d-s}{d-c}c+\frac{s-c}{d-c}d\right)dsdt
\end{aligned}$$

where

$$p(x,t)=\begin{cases}(t-a), & t\in\left[a,\frac{a+b}{2}\right]\\[2mm] (t-b), & t\in\left(\frac{a+b}{2},b\right]\end{cases}$$

and

$$q(y,s)=\begin{cases}(s-c), & s\in\left[c,\frac{c+d}{2}\right]\\[2mm] (s-d), & s\in\left(\frac{c+d}{2},d\right]\end{cases}.$$

for each $x\in[a,b]$ *and* $y\in[c,d]$.

Proof. Integration by parts, we can write

$$\begin{aligned}
&\frac{1}{(b-a)(d-c)}\int_a^b\int_c^d p(x,t)\,q(y,s)\,\frac{\partial^2 f}{\partial t\partial s}\left(\frac{b-t}{b-a}a+\frac{t-a}{b-a}b,\frac{d-s}{d-c}c+\frac{s-c}{d-c}d\right)dsdt \\
=\;&\int_c^d q(y,s)\left[\int_a^{\frac{a+b}{2}}(t-a)\,\frac{\partial^2 f}{\partial t\partial s}\left(\frac{b-t}{b-a}a+\frac{t-a}{b-a}b,\frac{d-s}{d-c}c+\frac{s-c}{d-c}d\right)dt\right. \\
&\left.+\int_{\frac{a+b}{2}}^b(t-b)\,\frac{\partial^2 f}{\partial t\partial s}\left(\frac{b-t}{b-a}a+\frac{t-a}{b-a}b,\frac{d-s}{d-c}c+\frac{s-c}{d-c}d\right)dt\right]ds
\end{aligned}$$

$$\begin{aligned}
=\;&\int_c^d q(y,s)\left\{\left[(t-a)\,\frac{\partial f}{\partial s}\left(\frac{b-t}{b-a}a+\frac{t-a}{b-a}b,\frac{d-s}{d-c}c+\frac{s-c}{d-c}d\right)\right]_a^{\frac{a+b}{2}}\right. \\
&-\int_a^{\frac{a+b}{2}}\frac{\partial f}{\partial s}\left(\frac{b-t}{b-a}a+\frac{t-a}{b-a}b,\frac{d-s}{d-c}c+\frac{s-c}{d-c}d\right)dt \\
&+\left[(t-b)\,\frac{\partial f}{\partial s}\left(\frac{b-t}{b-a}a+\frac{t-a}{b-a}b,\frac{d-s}{d-c}c+\frac{s-c}{d-c}d\right)\right]_{\frac{a+b}{2}}^b \\
&\left.-\int_{\frac{a+b}{2}}^b\frac{\partial f}{\partial s}\left(\frac{b-t}{b-a}a+\frac{t-a}{b-a}b,\frac{d-s}{d-c}c+\frac{s-c}{d-c}d\right)dt\right\}ds.
\end{aligned}$$

We obtain

$$\begin{aligned}
&\frac{1}{(b-a)(d-c)}\int_a^b\int_c^d p(x,t)\,q(y,s)\,\frac{\partial^2 f}{\partial t\partial s}\left(\frac{b-t}{b-a}a+\frac{t-a}{b-a}b,\frac{d-s}{d-c}c+\frac{s-c}{d-c}d\right)dsdt \\
=\;&(b-a)\int_c^d q(y,s)\left\{\frac{\partial f}{\partial s}\left(\frac{a+b}{2},\frac{d-s}{d-c}c+\frac{s-c}{d-c}d\right)\right. \\
&\left.-\int_a^b\frac{\partial f}{\partial s}\left(\frac{b-t}{b-a}a+\frac{t-a}{b-a}b,\frac{d-s}{d-c}c+\frac{s-c}{d-c}d\right)dt\right\}ds.
\end{aligned}$$

By integrating again, we get

$$\begin{aligned}
&\frac{1}{(b-a)(d-c)}\int_a^b\int_c^d p(x,t)\,q(y,s)\,\frac{\partial^2 f}{\partial t\partial s}\left(\frac{b-t}{b-a}a+\frac{t-a}{b-a}b,\frac{d-s}{d-c}c+\frac{s-c}{d-c}d\right)dsdt \\
=\;&(b-a)\left\{\int_c^{\frac{c+d}{2}}(s-c)\,\frac{\partial f}{\partial s}\left(\frac{a+b}{2},\frac{d-s}{d-c}c+\frac{s-c}{d-c}d\right)ds\right. \\
&+\int_{\frac{c+d}{2}}^d(s-d)\,\frac{\partial f}{\partial s}\left(\frac{a+b}{2},\frac{d-s}{d-c}c+\frac{s-c}{d-c}d\right)ds \\
&-\int_a^b\left[\int_c^{\frac{c+d}{2}}(s-c)\,\frac{\partial f}{\partial s}\left(\frac{b-t}{b-a}a+\frac{t-a}{b-a}b,\frac{d-s}{d-c}c+\frac{s-c}{d-c}d\right)ds\right. \\
&\left.\left.+\int_{\frac{c+d}{2}}^d(s-d)\,\frac{\partial f}{\partial s}\left(\frac{b-t}{b-a}a+\frac{t-a}{b-a}b,\frac{d-s}{d-c}c+\frac{s-c}{d-c}d\right)ds\right]dt\right\}.
\end{aligned}$$

By calculating the above integrals, we have

$$\begin{aligned}
& \frac{1}{(b-a)(d-c)}\int_a^b\int_c^d p(x,t)\,q(y,s)\frac{\partial^2 f}{\partial t\partial s}\left(\frac{b-t}{b-a}a+\frac{t-a}{b-a}b,\frac{d-s}{d-c}c+\frac{s-c}{d-c}d\right)dsdt \\
= & (b-a)(d-c)\,f\left(\frac{a+b}{2},\frac{c+d}{2}\right) \\
& -(b-a)\int_c^d f\left(\frac{a+b}{2},\frac{d-s}{d-c}c+\frac{s-c}{d-c}d\right)ds \\
& -(d-c)\int_a^b f\left(\frac{b-t}{b-a}a+\frac{t-a}{b-a}b,\frac{c+d}{2}\right)dt \\
& \int_a^b\int_c^d f\left(\frac{b-t}{b-a}a+\frac{t-a}{b-a}b,\frac{d-s}{d-c}c+\frac{s-c}{d-c}d\right)dsdt.
\end{aligned}$$

Using the change of the variable $x=\frac{b-t}{b-a}a+\frac{t-a}{b-a}b$ and $y=\frac{d-s}{d-c}c+\frac{s-c}{d-c}d$, then dividing both sides with $(b-a)\times(d-c)$, this completes the proof. □

Theorem 7. *Let $f:\Delta=[a,b]\times[c,d]\to\mathbb{R}$ be a partial differentiable mapping on $\Delta=[a,b]\times[c,d]$. If $\left|\frac{\partial^2 f}{\partial t\partial s}\right|$ is a convex function on the co-ordinates on Δ, then the following inequality holds;*

$$\begin{aligned}
& \left|f\left(\frac{a+b}{2},\frac{c+d}{2}\right)\right. \\
& -\frac{1}{(d-c)}\int_c^d f\left(\frac{a+b}{2},y\right)dy-\frac{1}{(b-a)}\int_a^b f\left(x,\frac{c+d}{2}\right)dx \\
& \left.+\frac{1}{(b-a)(d-c)}\int_a^b\int_c^d f(x,y)\,dydx\right| \\
\leq & \frac{(b-a)(d-c)}{64}\left[\left|\frac{\partial^2 f}{\partial t\partial s}(a,c)\right|+\left|\frac{\partial^2 f}{\partial t\partial s}(b,c)\right|+\left|\frac{\partial^2 f}{\partial t\partial s}(a,d)\right|+\left|\frac{\partial^2 f}{\partial t\partial s}(b,d)\right|\right].
\end{aligned} \tag{2.1}$$

Proof. From Lemma 2 and using the property of modulus, we have

$$\begin{aligned}
& \left|f\left(\frac{a+b}{2},\frac{c+d}{2}\right)\right. \\
& -\frac{1}{(d-c)}\int_c^d f\left(\frac{a+b}{2},y\right)dy-\frac{1}{(b-a)}\int_a^b f\left(x,\frac{c+d}{2}\right)dx \\
& \left.+\frac{1}{(b-a)(d-c)}\int_a^b\int_c^d f(x,y)\,dydx\right| \\
\leq & \frac{1}{(b-a)(d-c)}\int_a^b\int_c^d |p(x,t)\,q(y,s)|\left|\frac{\partial^2 f}{\partial t\partial s}\left(\frac{b-t}{b-a}a+\frac{t-a}{b-a}b,\frac{d-s}{d-c}c+\frac{s-c}{d-c}d\right)\right|dsdt
\end{aligned}$$

Since $\left|\frac{\partial^2 f}{\partial t \partial s}\right|$ is co-ordinated convex, we can write

$$\begin{aligned}
&\left| f\left(\frac{a+b}{2},\frac{c+d}{2}\right)\right. \\
&-\frac{1}{(d-c)}\int_c^d f\left(\frac{a+b}{2},y\right)dy-\frac{1}{(b-a)}\int_a^b f\left(x,\frac{c+d}{2}\right)dx \\
&\left.+\frac{1}{(b-a)(d-c)}\int_a^b\int_c^d f(x,y)\,dydx\right| \\
\leq\ & \frac{1}{(b-a)(d-c)} \\
&\times\int_c^d |q(y,s)|\left\{\int_a^{\frac{a+b}{2}}(t-a)\left[\frac{b-t}{b-a}\left|\frac{\partial^2 f}{\partial t\partial s}\left(a,\frac{d-s}{d-c}c+\frac{s-c}{d-c}d\right)\right|\right]dt\right. \\
&+\int_a^{\frac{a+b}{2}}(t-a)\left[\frac{t-a}{b-a}\left|\frac{\partial^2 f}{\partial t\partial s}\left(b,\frac{d-s}{d-c}c+\frac{s-c}{d-c}d\right)\right|\right]dt \\
&+\int_{\frac{a+b}{2}}^b(b-t)\left[\frac{b-t}{b-a}\left|\frac{\partial^2 f}{\partial t\partial s}\left(a,\frac{d-s}{d-c}c+\frac{s-c}{d-c}d\right)\right|\right]dt \\
&\left.+\int_{\frac{a+b}{2}}^b(b-t)\left[\frac{t-a}{b-a}\left|\frac{\partial^2 f}{\partial t\partial s}\left(b,\frac{d-s}{d-c}c+\frac{s-c}{d-c}d\right)\right|\right]dt\right\}ds.
\end{aligned}$$

By computing these integrals, we obtain

$$\begin{aligned}
&\left| f\left(\frac{a+b}{2},\frac{c+d}{2}\right)\right. \\
&-\frac{1}{(d-c)}\int_c^d f\left(\frac{a+b}{2},y\right)ds-\frac{1}{(b-a)}\int_a^b f\left(x,\frac{c+d}{2}\right)dx \\
&\left.+\frac{1}{(b-a)(d-c)}\int_a^b\int_c^d f(x,y)\,dxdy\right| \\
\leq\ & \frac{(b-a)}{8(d-c)}\left[\int_c^d |q(y,s)|\left|\frac{\partial^2 f}{\partial t\partial s}\left(a,\frac{d-s}{d-c}c+\frac{s-c}{d-c}d\right)\right|\right. \\
&\left.+\int_c^d |q(y,s)|\left|\frac{\partial^2 f}{\partial t\partial s}\left(b,\frac{d-s}{d-c}c+\frac{s-c}{d-c}d\right)\right|\right]ds.
\end{aligned}$$

Using co-ordinated convexity of $\left|\frac{\partial^2 f}{\partial t\partial s}\right|$ again, we get

$$\begin{aligned}
&\left|f\left(\frac{a+b}{2},\frac{c+d}{2}\right)\right.\\
&-\frac{1}{(d-c)}\int_c^d f\left(\frac{a+b}{2},y\right)dy-\frac{1}{(b-a)}\int_a^b f\left(x,\frac{c+d}{2}\right)dx\\
&\left.+\frac{1}{(b-a)(d-c)}\int_a^b\int_c^d f(x,y)\,dydx\right|\\
\leq\;&\frac{(b-a)}{8(d-c)}\left[\int_c^{\frac{c+d}{2}}(s-c)\left[\frac{d-s}{d-c}\left|\frac{\partial^2 f}{\partial t\partial s}(a,c)\right|\right]ds+\int_c^{\frac{c+d}{2}}(s-c)\left[\frac{s-c}{d-c}\left|\frac{\partial^2 f}{\partial t\partial s}(a,d)\right|\right]ds\right.\\
&+\int_{\frac{c+d}{2}}^d(d-s)\left[\frac{d-s}{d-c}\left|\frac{\partial^2 f}{\partial t\partial s}(a,c)\right|\right]ds+\int_{\frac{c+d}{2}}^d(d-s)\left[\frac{s-c}{d-c}\left|\frac{\partial^2 f}{\partial t\partial s}(a,d)\right|\right]ds\\
&+\int_c^{\frac{c+d}{2}}(s-c)\left[\frac{d-s}{d-c}\left|\frac{\partial^2 f}{\partial t\partial s}(b,c)\right|\right]ds+\int_c^{\frac{c+d}{2}}(s-c)\left[\frac{s-c}{d-c}\left|\frac{\partial^2 f}{\partial t\partial s}(b,d)\right|\right]ds\\
&\left.+\int_{\frac{c+d}{2}}^d(d-s)\left[\frac{d-s}{d-c}\left|\frac{\partial^2 f}{\partial t\partial s}(b,c)\right|\right]ds+\int_{\frac{c+d}{2}}^d(d-s)\left[\frac{s-c}{d-c}\left|\frac{\partial^2 f}{\partial t\partial s}(b,d)\right|\right]ds\right\}.
\end{aligned}$$

By a simple computation, we get the required result. □

Remark 1. *Suppose that all the assumptions of Theorem 7 are satisfied. If we choose* $\frac{\partial^2 f}{\partial t\partial s}$ *is bounded, i.e.,*

$$\left\|\frac{\partial^2 f(t,s)}{\partial t\partial s}\right\|_\infty=\sup_{(t,s)\in(a,b)\times(c,d)}\left|\frac{\partial^2 f(t,s)}{\partial t\partial s}\right|<\infty,$$

we get

$$\begin{aligned}
&\left|f\left(\frac{a+b}{2},\frac{c+d}{2}\right)\right.\\
&-\frac{1}{(d-c)}\int_c^d f\left(\frac{a+b}{2},y\right)dy-\frac{1}{(b-a)}\int_a^b f\left(x,\frac{c+d}{2}\right)dx\\
&\left.+\frac{1}{(b-a)(d-c)}\int_a^b\int_c^d f(x,y)\,dydx\right|\\
\leq\;&\frac{(b-a)(d-c)}{16}\left\|\frac{\partial^2 f(t,s)}{\partial t\partial s}\right\|_\infty
\end{aligned}\tag{2.2}$$

which is the inequality (*1.7*).

Theorem 8. *Let* $f:\Delta=[a,b]\times[c,d]\to\mathbb{R}$ *be a partial differentiable mapping on* $\Delta=[a,b]\times[c,d]$. *If* $\left|\frac{\partial^2 f}{\partial t\partial s}\right|^q$, $q>1$, *is a convex function on the co-ordinates on* Δ, *then the following inequality holds;*

$$
\begin{aligned}
&\left|f\left(\frac{a+b}{2},\frac{c+d}{2}\right)+\frac{1}{(b-a)(d-c)}\int_a^b\int_c^d f(x,y)\,dydx\right. \\
&\left.-\frac{1}{(d-c)}\int_c^d f\left(\frac{a+b}{2},y\right)dy-\frac{1}{(b-a)}\int_a^b f\left(x,\frac{c+d}{2}\right)dx\right| \\
\leq\ &\frac{(b-a)(d-c)}{4(p+1)^{\frac{2}{p}}} \\
&\times\left(\frac{\left|\frac{\partial^2 f}{\partial t\partial s}(a,c)\right|^q+\left|\frac{\partial^2 f}{\partial t\partial s}(b,c)\right|^q+\left|\frac{\partial^2 f}{\partial t\partial s}(a,d)\right|^q+\left|\frac{\partial^2 f}{\partial t\partial s}(b,d)\right|^q}{4}\right)^{\frac{1}{q}}.
\end{aligned}
\tag{2.3}
$$

Proof. From Lemma 2, we have

$$
\begin{aligned}
&\left|f\left(\frac{a+b}{2},\frac{c+d}{2}\right)+\frac{1}{(b-a)(d-c)}\int_a^b\int_c^d f(x,y)\,dydx\right. \\
&\left.-\frac{1}{(d-c)}\int_c^d f\left(\frac{a+b}{2},y\right)dy-\frac{1}{(b-a)}\int_a^b f\left(x,\frac{c+d}{2}\right)dx\right| \\
\leq\ &\frac{1}{(b-a)(d-c)}\int_a^b\int_c^d |p(x,t)\,q(y,s)|\left|\frac{\partial^2 f}{\partial t\partial s}\left(\frac{b-t}{b-a}a+\frac{t-a}{b-a}b,\frac{d-s}{d-c}c+\frac{s-c}{d-c}d\right)\right|dsdt
\end{aligned}
$$

By applying the well-known Hölder inequality for double integrals, then one has

$$
\begin{aligned}
&\left|f\left(\frac{a+b}{2},\frac{c+d}{2}\right)+\frac{1}{(b-a)(d-c)}\int_a^b\int_c^d f(x,y)\,dydx\right. \\
&\left.-\frac{1}{(d-c)}\int_c^d f\left(\frac{a+b}{2},y\right)dy-\frac{1}{(b-a)}\int_a^b f\left(x,\frac{c+d}{2}\right)dx\right| \\
\leq\ &\frac{1}{(b-a)(d-c)}\left\{\left(\int_a^b\int_c^d |p(x,t)\,q(y,s)|^p\,dtds\right)^{\frac{1}{p}}\right. \\
&\left.\times\left(\int_a^b\int_c^d\left|\frac{\partial^2 f}{\partial t\partial s}\left(\frac{b-t}{b-a}a+\frac{t-a}{b-a}b,\frac{d-s}{d-c}c+\frac{s-c}{d-c}d\right)\right|^q dsdt\right)^{\frac{1}{q}}\right\}.
\end{aligned}
\tag{2.4}
$$

Since $\left|\frac{\partial^2 f}{\partial t\partial s}\right|^q$ is a co-ordinated convex function on Δ, we can write for all $(t,s)\in[a,b]\times[c,d]$

$$
\begin{aligned}
&\left|\frac{\partial^2 f}{\partial t\partial s}\left(\frac{b-t}{b-a}a+\frac{t-a}{b-a}b,\frac{d-s}{d-c}c+\frac{s-c}{d-c}d\right)\right|^q \\
\leq\ &\frac{b-t}{b-a}\left|\frac{\partial^2 f}{\partial t\partial s}\left(a,\frac{d-s}{d-c}c+\frac{s-c}{d-c}d\right)\right|^q \\
&+\frac{t-a}{b-a}\left|\frac{\partial^2 f}{\partial t\partial s}\left(b,\frac{d-s}{d-c}c+\frac{s-c}{d-c}d\right)\right|^q
\end{aligned}
$$

and

$$\begin{aligned}
&\left|\frac{\partial^2 f}{\partial t\partial s}\left(\frac{b-t}{b-a}a+\frac{t-a}{b-a}b,\frac{d-s}{d-c}c+\frac{s-c}{d-c}d\right)\right|^q \qquad (2.5)\\
\leq\ & \left(\frac{b-t}{b-a}\right)\left(\frac{d-s}{d-c}\right)\left|\frac{\partial^2 f}{\partial t\partial s}(a,c)\right|^q\\
&+\left(\frac{b-t}{b-a}\right)\left(\frac{s-c}{d-c}\right)\left|\frac{\partial^2 f}{\partial t\partial s}(a,d)\right|^q\\
&+\left(\frac{t-a}{b-a}\right)\left(\frac{d-s}{d-c}\right)\left|\frac{\partial^2 f}{\partial t\partial s}(b,c)\right|^q\\
&+\left(\frac{t-a}{b-a}\right)\left(\frac{s-c}{d-c}\right)\left|\frac{\partial^2 f}{\partial t\partial s}(b,d)\right|^q.
\end{aligned}$$

Using inequality of (2.5) in (2.4), we get

$$\begin{aligned}
&\Bigg|f\left(\frac{a+b}{2},\frac{c+d}{2}\right)+\frac{1}{(b-a)(d-c)}\int_a^b\int_c^d f(x,y)\,dydx\\
&-\frac{1}{(d-c)}\int_c^d f\left(\frac{a+b}{2},y\right)dy-\frac{1}{(b-a)}\int_a^b f\left(x,\frac{c+d}{2}\right)dx\Bigg|\\
\leq\ & \frac{(b-a)(d-c)}{4(p+1)^{\frac{2}{p}}}\left(\frac{\left|\frac{\partial^2 f}{\partial t\partial s}(a,c)\right|^q+\left|\frac{\partial^2 f}{\partial t\partial s}(b,c)\right|^q+\left|\frac{\partial^2 f}{\partial t\partial s}(a,d)\right|^q+\left|\frac{\partial^2 f}{\partial t\partial s}(b,d)\right|^q}{4}\right)^{\frac{1}{q}}
\end{aligned}$$

where we have used the fact that

$$\left(\int_a^b\int_c^d |p(x,t)\,q(y,s)|^p\,dtds\right)^{\frac{1}{p}}=\frac{[(b-a)(d-c)]^{1+\frac{1}{p}}}{4(p+1)^{\frac{2}{p}}}.$$

This completes the proof. □

Remark 2. *Suppose that all the assumptions of Theorem 8 are satisfied. If we choose* $\frac{\partial^2 f}{\partial t\partial s}$ *is bounded, i.e.,*

$$\left\|\frac{\partial^2 f(t,s)}{\partial t\partial s}\right\|_\infty=\sup_{(t,s)\in(a,b)\times(c,d)}\left|\frac{\partial^2 f(t,s)}{\partial t\partial s}\right|<\infty,$$

we get

$$\begin{aligned}
&\Bigg|f\left(\frac{a+b}{2},\frac{c+d}{2}\right) \qquad (2.6)\\
&-\frac{1}{(d-c)}\int_c^d f\left(\frac{a+b}{2},y\right)dy-\frac{1}{(b-a)}\int_a^b f\left(x,\frac{c+d}{2}\right)dx\\
&+\frac{1}{(b-a)(d-c)}\int_a^b\int_c^d f(x,y)\,dydx\Bigg|\\
\leq\ & \frac{(b-a)(d-c)}{4(p+1)^{\frac{2}{p}}}\left\|\frac{\partial^2 f(t,s)}{\partial t\partial s}\right\|_\infty.
\end{aligned}$$

Theorem 9. *Let* $f:\Delta=[a,b]\times[c,d]\to\mathbb{R}$ *be a partial differentiable mapping on* $\Delta=[a,b]\times[c,d]$. *If* $\left|\frac{\partial^2 f}{\partial t\partial s}\right|^q$, $q\geq 1$, *is a convex function on the co-ordinates on* Δ, *then the following inequality holds;*

$$\begin{aligned}
&\left|f\left(\frac{a+b}{2},\frac{c+d}{2}\right)+\frac{1}{(b-a)(d-c)}\int_a^b\int_c^d f(x,y)\,dydx\right.\\
&\left.-\frac{1}{(d-c)}\int_c^d f\left(\frac{a+b}{2},y\right)dy-\frac{1}{(b-a)}\int_a^b f\left(x,\frac{c+d}{2}\right)dx\right|\\
\leq\ &\frac{(b-a)(d-c)}{16}\\
&\times\left(\frac{\left|\frac{\partial^2 f}{\partial t\partial s}(a,c)\right|^q+\left|\frac{\partial^2 f}{\partial t\partial s}(b,c)\right|^q+\left|\frac{\partial^2 f}{\partial t\partial s}(a,d)\right|^q+\left|\frac{\partial^2 f}{\partial t\partial s}(b,d)\right|^q}{4}\right)^{\frac{1}{q}}.
\end{aligned}\tag{2.7}$$

Proof. From Lemma 2, we have

$$\begin{aligned}
&\left|f\left(\frac{a+b}{2},\frac{c+d}{2}\right)+\frac{1}{(b-a)(d-c)}\int_a^b\int_c^d f(x,y)\,dydx\right.\\
&\left.-\frac{1}{(d-c)}\int_c^d f\left(\frac{a+b}{2},y\right)dy-\frac{1}{(b-a)}\int_a^b f\left(x,\frac{c+d}{2}\right)dx\right|\\
\leq\ &\frac{1}{(b-a)(d-c)}\\
&\int_a^b\int_c^d |p(x,t)\,q(y,s)|\left|\frac{\partial^2 f}{\partial t\partial s}\left(\frac{b-t}{b-a}a+\frac{t-a}{b-a}b,\frac{d-s}{d-c}c+\frac{s-c}{d-c}d\right)\right|dsdt
\end{aligned}$$

By applying the well-known Power mean inequality for double integrals, then one has

$$\begin{aligned}
&\left|f\left(\frac{a+b}{2},\frac{c+d}{2}\right)+\frac{1}{(b-a)(d-c)}\int_a^b\int_c^d f(x,y)\,dydx\right.\\
&\left.-\frac{1}{(d-c)}\int_c^d f\left(\frac{a+b}{2},y\right)dy-\frac{1}{(b-a)}\int_a^b f\left(x,\frac{c+d}{2}\right)dx\right|\\
\leq\ &\frac{1}{(b-a)(d-c)}\times\left(\int_a^b\int_c^d |p(x,t)\,q(y,s)|\,dsdt\right)^{1-\frac{1}{q}}\\
&\left(\int\limits_a^b\int\limits_c^d |p(x,t)\,q(y,s)|\left|\frac{\partial^2 f}{\partial t\partial s}\left(\frac{b-t}{b-a}a+\frac{t-a}{b-a}b,\frac{d-s}{d-c}c+\frac{s-c}{d-c}d\right)\right|^q dsdt\right)^{\frac{1}{q}}.
\end{aligned}\tag{2.8}$$

Since $\left|\frac{\partial^2 f}{\partial t\partial s}\right|^q$ is a co-ordinated convex function on Δ, we can write for all $(t,s) \in [a,b]\times[c,d]$

$$\begin{aligned}
&\left|\frac{\partial^2 f}{\partial t\partial s}\left(\frac{b-t}{b-a}a+\frac{t-a}{b-a}b,\frac{d-s}{d-c}c+\frac{s-c}{d-c}d\right)\right|^q \\
\leq\ & \left(\frac{b-t}{b-a}\right)\left(\frac{d-s}{d-c}\right)\left|\frac{\partial^2 f}{\partial t\partial s}(a,c)\right|^q \\
&+\left(\frac{b-t}{b-a}\right)\left(\frac{s-c}{d-c}\right)\left|\frac{\partial^2 f}{\partial t\partial s}(a,d)\right|^q \\
&+\left(\frac{t-a}{b-a}\right)\left(\frac{d-s}{d-c}\right)\left|\frac{\partial^2 f}{\partial t\partial s}(b,c)\right|^q \\
&+\left(\frac{t-a}{b-a}\right)\left(\frac{s-c}{d-c}\right)\left|\frac{\partial^2 f}{\partial t\partial s}(b,d)\right|^q .
\end{aligned} \tag{2.9}$$

If we use (2.9) in (2.8), we get

$$\begin{aligned}
&\left|f\left(\frac{a+b}{2},\frac{c+d}{2}\right)+\frac{1}{(b-a)(d-c)}\int_a^b\int_c^d f(x,y)\,dydx\right. \\
&\left.-\frac{1}{(d-c)}\int_c^d f\left(\frac{a+b}{2},y\right)dy-\frac{1}{(b-a)}\int_a^b f\left(x,\frac{c+d}{2}\right)dx\right| \\
\leq\ & \frac{1}{(b-a)(d-c)}\left\{\left(\int_a^b\int_c^d |p(x,t)\,q(y,s)|\,dsdt\right)^{1-\frac{1}{q}}\right. \\
&\times\left(\int_a^b\int_c^d |p(x,t)\,q(y,s)|\left|\frac{\partial^2 f}{\partial t\partial s}\left(\frac{b-t}{b-a}a+\frac{t-a}{b-a}b,\frac{d-s}{d-c}c+\frac{s-c}{d-c}d\right)\right|^q dsdt\right)^{\frac{1}{q}}
\end{aligned}$$

hence it follows that

$$\begin{aligned}
&\left|f\left(\frac{a+b}{2},\frac{c+d}{2}\right)+\frac{1}{(b-a)(d-c)}\int_a^b\int_c^d f(x,y)\,dydx\right. \\
&\left.-\frac{1}{(d-c)}\int_c^d f\left(\frac{a+b}{2},y\right)dy-\frac{1}{(b-a)}\int_a^b f\left(x,\frac{c+d}{2}\right)dx\right| \\
\leq\ & \frac{1}{(b-a)(d-c)}\left\{\left(\int_a^b\int_c^d |p(x,t)\,q(y,s)|\,dsdt\right)^{1-\frac{1}{q}}\right. \\
&\times\left(\int_a^b\int_c^d |p(x,t)\,q(y,s)|\left[\left(\frac{b-t}{b-a}\right)\left(\frac{d-s}{d-c}\right)\left|\frac{\partial^2 f}{\partial t\partial s}(a,c)\right|^q+\left(\frac{b-t}{b-a}\right)\left(\frac{s-c}{d-c}\right)\left|\frac{\partial^2 f}{\partial t\partial s}(a,d)\right|^q\right.\right. \\
&\left.\left.+\left(\frac{t-a}{b-a}\right)\left(\frac{d-s}{d-c}\right)\left|\frac{\partial^2 f}{\partial t\partial s}(b,c)\right|^q+\left(\frac{t-a}{b-a}\right)\left(\frac{s-c}{d-c}\right)\left|\frac{\partial^2 f}{\partial t\partial s}(b,d)\right|^q\right]\right)^{\frac{1}{q}} .
\end{aligned}$$

Computing the above integrals and using the fact that

$$\left(\int_a^b\int_c^d |p(x,t)\,q(y,s)|\,dtds\right)^{1-\frac{1}{q}}=\left(\frac{(b-a)^2(d-c)^2}{16}\right)^{1-\frac{1}{q}} .$$

This completes the proof. □

3. INEQUALITIES FOR CO-ORDINATED $s-$CONVEX FUNCTIONS

To prove our main results we need the following lemma:

Lemma 2. *Let* $f:\Delta\subset\mathbb{R}^2\to\mathbb{R}$ *be an absolutely continuous function on* Δ *where* $a<b$, $c<d$ *and* $t,\lambda\in[0,1]$, *if* $\frac{\partial^2 f}{\partial t\partial\lambda}\in L(\Delta)$, *then the following equality holds:*

$$\begin{aligned}
&\frac{f(a,c)+r_2 f(a,d)+r_1 f(b,c)+r_1 r_2 f(b,d)}{(r_1+1)(r_2+1)}\\
&+\frac{1}{(b-a)(d-c)}\int_a^b\int_c^d f(x,y)dxdy\\
&-\left(\frac{r_2}{r_2+1}\right)\frac{1}{d-c}\int_c^d f(b,y)dy-\left(\frac{1}{r_1+1}\right)\frac{1}{d-c}\int_c^d f(a,y)dy\\
&-\left(\frac{r_2}{r_2+1}\right)\frac{1}{b-a}\int_a^b f(x,d)dx-\left(\frac{1}{r_2+1}\right)\frac{1}{b-a}\int_a^b f(x,c)dx\\
=\;&\frac{(b-a)(d-c)}{(r_1+1)(r_2+1)}\\
&\times\int_0^1\int_0^1((r_1+1)t-1)((r_2+1)\lambda-1)\frac{\partial^2 f}{\partial t\partial\lambda}(tb+(1-t)a,\lambda d+(1-\lambda)c)\,dtd\lambda
\end{aligned}$$

for some fixed $r_1,r_2\in[0,1]$.

Proof. Integration by parts, we get

$$\begin{aligned}
&\int_0^1\int_0^1((r_1+1)t-1)((r_2+1)\lambda-1)\frac{\partial^2 f}{\partial t\partial\lambda}(tb+(1-t)a,\lambda d+(1-\lambda)c)\,dtd\lambda\\
=\;&\int_0^1((r_2+1)\lambda-1)\left[\int_0^1((r_1+1)t-1)\frac{\partial^2 f}{\partial t\partial\lambda}(tb+(1-t)a,\lambda d+(1-\lambda)c)\,dtd\lambda\right]\\
=\;&\int_0^1((r_2+1)\lambda-1)\left[\frac{((r_1+1)t-1)}{(b-a)}\frac{\partial f}{\partial\lambda}(tb+(1-t)a,\lambda d+(1-\lambda)c)\Big|_0^1\right.\\
&\left.-\frac{r_1+1}{b-a}\int_0^1\frac{\partial f}{\partial\lambda}(tb+(1-t)a,\lambda d+(1-\lambda)c)\,dt\right]d\lambda
\end{aligned}$$

$$\begin{aligned}
&\int_0^1\int_0^1((r_1+1)t-1)((r_2+1)\lambda-1)\frac{\partial^2 f}{\partial t\partial\lambda}(tb+(1-t)a,\lambda d+(1-\lambda)c)\,dtd\lambda\\
=\;&\int_0^1((r_2+1)\lambda-1)\left[\frac{r_1}{b-a}\frac{\partial f}{\partial\lambda}(b,\lambda d+(1-\lambda)c)+\frac{1}{b-a}\frac{\partial f}{\partial\lambda}(a,\lambda d+(1-\lambda)c)\right.\\
&\left.-\frac{r_1+1}{b-a}\int_0^1\frac{\partial f}{\partial\lambda}(tb+(1-t)a,\lambda d+(1-\lambda)c)\,dt\right]d\lambda
\end{aligned}$$

Again by integration by parts, we have

$$\begin{aligned}
&\int_0^1 ((r_2+1)\lambda-1)\left[\frac{r_1}{b-a}\frac{\partial f}{\partial\lambda}(b,\lambda d+(1-\lambda)c)+\frac{1}{b-a}\frac{\partial f}{\partial\lambda}(a,\lambda d+(1-\lambda)c)\right.\\
&\left.-\frac{r_1+1}{b-a}\int_0^1\frac{\partial f}{\partial\lambda}(tb+(1-t)a,\lambda d+(1-\lambda)c)\,dt\right]d\lambda\\
= \quad &\frac{r_1}{b-a}\frac{((r_2+1)\lambda-1)}{d-c}f(b,\lambda d+(1-\lambda)c)\Big|_0^1-\frac{r_1(r_2+1)}{(b-a)(d-c)}\int_0^1 f(b,\lambda d+(1-\lambda)c)\,d\lambda\\
&+\frac{1}{b-a}\frac{((r_2+1)\lambda-1)}{d-c}f(a,\lambda d+(1-\lambda)c)\Big|_0^1-\frac{(r_2+1)}{(b-a)(d-c)}\int_0^1 f(a,\lambda d+(1-\lambda)c)\,d\lambda\\
&-\frac{r_1+1}{b-a}\int_0^1\left[\int_0^1((r_2+1)\lambda-1)\frac{\partial f}{\partial\lambda}(tb+(1-t)a,\lambda d+(1-\lambda)c)\,d\lambda\right]dt.
\end{aligned}$$

Computing these integrals and by using the results, we obtain

$$\begin{aligned}
&\int_0^1\int_0^1((r_1+1)t-1)((r_2+1)\lambda-1)\frac{\partial^2 f}{\partial t\partial\lambda}(tb+(1-t)a,\lambda d+(1-\lambda)c)\,dtd\lambda\\
= \quad &\frac{1}{(b-a)(d-c)}\left[f(a,c)+r_2f(a,d)+r_1f(b,c)+r_1r_2f(b,d)\right.\\
&-r_1(r_2+1)\int_0^1 f(b,\lambda d+(1-\lambda)c)d\lambda-(r_2+1)\int_0^1 f(a,\lambda d+(1-\lambda)c)d\lambda\\
&-r_2(r_1+1)\int_0^1 f(tb+(1-t)a,d)dt-(r_2+1)\int_0^1 f(tb+(1-t)a,c)dt\\
&\left.(r_1+1)(r_2+1)\int_0^1\int_0^1 f(tb+(1-t)a,\lambda d+(1-\lambda)c)\,dtd\lambda\right].
\end{aligned}$$

Using the change of the variable $x=tb+(1-t)a$ and $y=\lambda d+(1-\lambda)c$ for $t,\lambda\in[0,1]$ and multiplying the both sides by $\frac{(b-a)(d-c)}{(r_1+1)(r_2+1)}$, we get the required result. □

Theorem 10. *Let* $f:\Delta=[a,b]\times[c,d]\subset[0,\infty)^2\to[0,\infty)$ *be an absolutely continuous function on* Δ. *If* $\left|\frac{\partial^2 f}{\partial t\partial\lambda}\right|$ *is* $s-$*convex function on the co-ordinates on*

Δ, *then one has the inequality:*

$$
\begin{aligned}
&\left|\frac{f(a,c)+r_2 f(a,d)+r_1 f(b,c)+r_1 r_2 f(b,d)}{(r_1+1)(r_2+1)}\right. \\
&-\frac{1}{d-c}\left[\left(\frac{r_2}{r_2+1}\right)\int_c^d f(b,y)dy+\left(\frac{1}{r_1+1}\right)\int_c^d f(a,y)dy\right] \\
&-\frac{1}{b-a}\left[\left(\frac{r_2}{r_2+1}\right)\int_a^b f(x,d)dx+\left(\frac{1}{r_2+1}\right)\int_a^b f(x,c)dx\right] \\
&\left.+\frac{1}{(b-a)(d-c)}\int_a^b\int_c^d f(x,y)dxdy\right| \\
\leq\ & \frac{(b-a)(d-c)}{(r_1+1)(r_2+1)(s+1)^2(s+2)^2} \\
&\times\left(MN\left|\frac{\partial^2 f}{\partial t\partial\lambda}\right|(a,c)+LN\left|\frac{\partial^2 f}{\partial t\partial\lambda}\right|(a,d)\right. \\
&\left.+KM\left|\frac{\partial^2 f}{\partial t\partial\lambda}\right|(b,c)+KL\left|\frac{\partial^2 f}{\partial t\partial\lambda}\right|(b,d)\right)
\end{aligned} \tag{3.1}
$$

where

$$
\begin{aligned}
M &= \left(s+1+2(r_1+1)\left(\frac{r_1}{r_1+1}\right)^{s+2}-r_1\right) \\
N &= \left(s+1+2(r_2+1)\left(\frac{r_2}{r_2+1}\right)^{s+2}-r_2\right) \\
L &= \left(r_2(s+1)+2\left(\frac{1}{r_2+1}\right)^{s+1}-1\right) \\
K &= \left(r_1(s+1)+2\left(\frac{1}{r_1+1}\right)^{s+1}-1\right).
\end{aligned}
$$

Proof. From Lemma 3, we can write;

$$
\begin{aligned}
&\left|\frac{f(a,c)+r_2 f(a,d)+r_1 f(b,c)+r_1 r_2 f(b,d)}{(r_1+1)(r_2+1)}+\frac{1}{(b-a)(d-c)}\int_a^b\int_c^d f(x,y)dxdy\right. \\
&-\left(\frac{r_2}{r_2+1}\right)\frac{1}{d-c}\int_c^d f(b,y)dy-\left(\frac{1}{r_1+1}\right)\frac{1}{d-c}\int_c^d f(a,y)dy \\
&\left.-\left(\frac{r_2}{r_2+1}\right)\frac{1}{b-a}\int_a^b f(x,d)dx-\left(\frac{1}{r_2+1}\right)\frac{1}{b-a}\int_a^b f(x,c)dx\right| \\
=\ & \frac{(b-a)(d-c)}{(r_1+1)(r_2+1)} \\
&\times\int_0^1\int_0^1\left|((r_1+1)t-1)((r_2+1)\lambda-1)\right|\left|\frac{\partial^2 f}{\partial t\partial\lambda}(tb+(1-t)a,\lambda d+(1-\lambda)c)\right|dtd\lambda
\end{aligned}
$$

By using co-ordinated $s-$convexity of f, we have

$$\begin{aligned}
&\left|\frac{f\left(a,c\right)+r_2f\left(a,d\right)+r_1f\left(b,c\right)+r_1r_2f\left(b,d\right)}{\left(r_1+1\right)\left(r_2+1\right)}\right. \\
&-\left(\frac{r_2}{r_2+1}\right)\frac{1}{d-c}\int_c^d f(b,y)dy-\left(\frac{1}{r_1+1}\right)\frac{1}{d-c}\int_c^d f(a,y)dy \\
&-\left(\frac{r_2}{r_2+1}\right)\frac{1}{b-a}\int_a^b f(x,d)dx-\left(\frac{1}{r_2+1}\right)\frac{1}{b-a}\int_a^b f(x,c)dx \\
&\left.+\frac{1}{(b-a)(d-c)}\int_a^b\int_c^d f(x,y)dxdy\right| \\
= &\ \frac{(b-a)(d-c)}{\left(r_1+1\right)\left(r_2+1\right)} \\
&\times\int_0^1\left[\int_0^1\left|\left(\left(r_1+1\right)t-1\right)\left(\left(r_2+1\right)\lambda-1\right)\right|\right. \\
&\left.\left\{t^s\left|\frac{\partial^2 f}{\partial t\partial\lambda}\left(b,\lambda d+\left(1-\lambda\right)c\right)\right|+\left(1-t\right)^s\left|\frac{\partial^2 f}{\partial t\partial\lambda}\left(a,\lambda d+\left(1-\lambda\right)c\right)\right|\right\}dt\right]d\lambda.
\end{aligned}$$

By calculating the above integrals, we get

$$\begin{aligned}
(3.2)\quad &\int_0^1\left|\left(\left(r_1+1\right)t-1\right)\right|\left\{t^s\left|\frac{\partial^2 f}{\partial t\partial\lambda}\left(b,\lambda d+\left(1-\lambda\right)c\right)\right|\right. \\
&\left.+\left(1-t\right)^s\left|\frac{\partial^2 f}{\partial t\partial\lambda}\left(a,\lambda d+\left(1-\lambda\right)c\right)\right|\right\}dt \\
= &\int_0^{\frac{1}{r_1+1}}\left(1-\left(r_1+1\right)t\right)\left\{t^s\left|\frac{\partial^2 f}{\partial t\partial\lambda}\left(b,\lambda d+\left(1-\lambda\right)c\right)\right|\right. \\
&\left.+\left(1-t\right)^s\left|\frac{\partial^2 f}{\partial t\partial\lambda}\left(a,\lambda d+\left(1-\lambda\right)c\right)\right|\right\}dt \\
&+\int_{\frac{1}{r_1+1}}^1\left(\left(r_1+1\right)t-1\right)\left\{t^s\left|\frac{\partial^2 f}{\partial t\partial\lambda}\left(b,\lambda d+\left(1-\lambda\right)c\right)\right|\right. \\
&\left.+\left(1-t\right)^s\left|\frac{\partial^2 f}{\partial t\partial\lambda}\left(a,\lambda d+\left(1-\lambda\right)c\right)\right|\right\}dt \\
= &\ \frac{1}{\left(s+1\right)\left(s+2\right)}\left[\left(r_1\left(s+1\right)+2\left(\frac{1}{r_1+1}\right)^{s+1}-1\right)\left|\frac{\partial^2 f}{\partial t\partial\lambda}\left(b,\lambda d+\left(1-\lambda\right)c\right)\right|\right. \\
&\left.+\left(s+1+2\left(r_1+1\right)\left(\frac{r_1}{r_1+1}\right)^{s+2}-r_1\right)\left|\frac{\partial^2 f}{\partial t\partial\lambda}\left(a,\lambda d+\left(1-\lambda\right)c\right)\right|\right].
\end{aligned}$$

By a similar argument for other integrals, by using co-ordinated $s-$convexity of f, we get

$$\begin{aligned}
&\int_0^1 \left|\left(\left(r_2+1\right)t-1\right)\right| \left\{ \left|\frac{\partial^2 f}{\partial t \partial \lambda}\left(b, \lambda d+(1-\lambda)c\right)\right| + \left|\frac{\partial^2 f}{\partial t \partial \lambda}\left(a, \lambda d+(1-\lambda)c\right)\right| \right\} d\lambda \\
=\;& \int_0^{\frac{1}{r_2+1}} \left(1-\left(r_2+1\right)t\right) \left\{ \lambda^s \left|\frac{\partial^2 f}{\partial t \partial \lambda}(b,d)\right| + (1-\lambda)^s \left|\frac{\partial^2 f}{\partial t \partial \lambda}(b,c)\right| \right\} d\lambda \\
&+ \int_{\frac{1}{r_2+1}}^1 \left(\left(r_2+1\right)t-1\right) \left\{ \lambda^s \left|\frac{\partial^2 f}{\partial t \partial \lambda}(a,d)\right| + (1-\lambda)^s \left|\frac{\partial^2 f}{\partial t \partial \lambda}(a,c)\right| \right\} d\lambda \\
=\;& \frac{1}{(s+1)(s+2)}\left[r_2(s+1)+2\left(\frac{1}{r_2+1}\right)^{s+1}-1\right]\left|\frac{\partial^2 f}{\partial t \partial \lambda}(b,d)\right| \\
&+\frac{1}{(s+1)(s+2)}\left[r_2(s+1)+2\left(\frac{1}{r_2+1}\right)^{s+1}-1\right]\left|\frac{\partial^2 f}{\partial t \partial \lambda}(a,d)\right| \\
&+\frac{1}{(s+1)(s+2)}\left[s+1+2\left(r_2+1\right)\left(\frac{r_2}{r_2+1}\right)^{s+2}-r_1\right]\left|\frac{\partial^2 f}{\partial t \partial \lambda}(b,c)\right| \\
&+\frac{1}{(s+1)(s+2)}\left[s+1+2\left(r_1+1\right)\left(\frac{r_2}{r_2+1}\right)^{s+2}-r_1\right]\left|\frac{\partial^2 f}{\partial t \partial \lambda}(a,c)\right|.
\end{aligned}$$

By using these in (3.2), we obtain the inequality (3.1). □

Corollary 2. *(1) If we choose* $r_1 = r_2 = 1$ *in (3.1), we have*

$$\begin{aligned}
&\left|\frac{f(a,c)+f(a,d)+f(b,c)+f(b,d)}{4}\right. \\
&-\frac{1}{2}\left[\frac{1}{d-c}\int_c^d \left[f(b,y)+f(a,y)\right]dy\right]-\frac{1}{2}\left[\frac{1}{b-a}\int_a^b \left[f(x,d)+f(x,c)\right]dx\right] \\
&\left.+\frac{1}{(b-a)(d-c)}\int_a^b\int_c^d f(x,y)dxdy\right| \\
\leq\;& \frac{(b-a)(d-c)}{(s+1)^2(s+2)^2}\left(s+\frac{1}{2^s}\right)^2 \\
&\left(\left|\frac{\partial^2 f}{\partial t \partial \lambda}\right|(a,c)+\left|\frac{\partial^2 f}{\partial t \partial \lambda}\right|(a,d)+\left|\frac{\partial^2 f}{\partial t \partial \lambda}\right|(b,c)+\left|\frac{\partial^2 f}{\partial t \partial \lambda}\right|(b,d)\right)
\end{aligned} \tag{3.3}$$

(2) If we choose $r_1 = r_2 = 0$ *in (3.1), we have*

$$\begin{aligned}
&\left|f(a,c)-\frac{1}{d-c}\int_c^d f(a,y)dy-\frac{1}{b-a}\int_a^b f(x,c)dx\right. \\
&\left.+\frac{1}{(b-a)(d-c)}\int_a^b\int_c^d f(x,y)dxdy\right| \\
\leq\;& \frac{(b-a)(d-c)}{(s+1)^2(s+2)^2} \\
&\left((s+1)^2\left|\frac{\partial^2 f}{\partial t \partial \lambda}\right|(a,c)+(s+1)\left|\frac{\partial^2 f}{\partial t \partial \lambda}\right|(a,d)+(s+1)\left|\frac{\partial^2 f}{\partial t \partial \lambda}\right|(b,c)+\left|\frac{\partial^2 f}{\partial t \partial \lambda}\right|(b,d)\right)
\end{aligned}$$

Remark 3. *If we choose* $s = 1$ *in (3.3), we get an improvement for the inequality (1.2).*

Theorem 11. *Let* $f : \Delta = [a, b] \times [c, d] \subset [0, \infty)^2 \to [0, \infty)$ *be an absolutely continuous function on* Δ*. If* $\left|\frac{\partial^2 f}{\partial t \partial \lambda}\right|^{\frac{p}{p-1}}$ *is* $s-$*convex function on the co-ordinates on* Δ*, for some fixed* $s \in (0, 1]$ *and* $p > 1$*, then one has the inequality:*

$$
\begin{aligned}
(3.4)\qquad & \left|\frac{f(a,c) + r_2 f(a,d) + r_1 f(b,c) + r_1 r_2 f(b,d)}{(r_1+1)(r_2+1)}\right. \\
& -\left(\frac{r_2}{r_2+1}\right)\frac{1}{d-c}\int_c^d f(b,y)dy - \left(\frac{1}{r_1+1}\right)\frac{1}{d-c}\int_c^d f(a,y)dy \\
& -\left(\frac{r_2}{r_2+1}\right)\frac{1}{b-a}\int_a^b f(x,d)dx - \left(\frac{1}{r_2+1}\right)\frac{1}{b-a}\int_a^b f(x,c)dx \\
& \left. +\frac{1}{(b-a)(d-c)}\int_a^b\int_c^d f(x,y)dxdy\right| \\
= \;& \frac{(b-a)(d-c)}{(r_1+1)(r_2+1)}\left(\frac{\left(1 + r_1^{\frac{p+1}{p}}\right)\left(1 + r_2^{\frac{p+1}{p}}\right)}{(r_1+1)^{\frac{1}{p}}(r_2+1)^{\frac{1}{p}}(p+1)^{\frac{2}{p}}}\right) \\
& \times\left(\frac{\left|\frac{\partial^2 f}{\partial t\partial\lambda}\right|^q(a,c) + \left|\frac{\partial^2 f}{\partial t\partial\lambda}\right|^q(a,d) + \left|\frac{\partial^2 f}{\partial t\partial\lambda}\right|^q(b,c) + \left|\frac{\partial^2 f}{\partial t\partial\lambda}\right|^q(b,d)}{(s+1)^2}\right)^{\frac{1}{q}}
\end{aligned}
$$

for some fixed $r_1, r_2 \in [0, 1]$ *, where* $q = \frac{p}{p-1}$.

Proof. Let $p > 1$. From Lemma 3 and using the Hölder inequality for double integrals, we can write

$$
\begin{aligned}
& \left|\frac{f(a,c) + r_2 f(a,d) + r_1 f(b,c) + r_1 r_2 f(b,d)}{(r_1+1)(r_2+1)}\right. \\
& -\left(\frac{r_2}{r_2+1}\right)\frac{1}{d-c}\int_c^d f(b,y)dy - \left(\frac{1}{r_1+1}\right)\frac{1}{d-c}\int_c^d f(a,y)dy \\
& -\left(\frac{r_2}{r_2+1}\right)\frac{1}{b-a}\int_a^b f(x,d)dx - \left(\frac{1}{r_2+1}\right)\frac{1}{b-a}\int_a^b f(x,c)dx \\
& \left. +\frac{1}{(b-a)(d-c)}\int_a^b\int_c^d f(x,y)dxdy\right| \\
= \;& \frac{(b-a)(d-c)}{(r_1+1)(r_2+1)}\left(\int_0^1\int_0^1 \left|\left((r_1+1)t-1\right)\left((r_2+1)\lambda-1\right)\right|^p dtd\lambda\right)^{\frac{1}{p}} \\
& \times\left(\int_0^1\int_0^1\left|\frac{\partial^2 f}{\partial t\partial\lambda}\left(tb + (1-t)a, \lambda d + (1-\lambda)c\right)\right|^q dtd\lambda\right)^{\frac{1}{q}}.
\end{aligned}
$$

Since $\left|\frac{\partial^2 f}{\partial t \partial \lambda}\right|^q$ is $s-$convex function on the co-ordinates on Δ, we can write for $t, \lambda \in [0,1]$

$$\begin{aligned}
&\left|\frac{\partial^2 f}{\partial t \partial \lambda}\left(tb+(1-t)a, \lambda d+(1-\lambda)c\right)\right|^q \\
\leq\ & t^s\left|\frac{\partial^2 f}{\partial t \partial \lambda}\left(b, \lambda d+(1-\lambda)c\right)\right|^q+(1-t)^s\left|\frac{\partial^2 f}{\partial t \partial \lambda}\left(a, \lambda d+(1-\lambda)c\right)\right|^q
\end{aligned}$$

and

$$\begin{aligned}
&\left|\frac{\partial^2 f}{\partial t \partial \lambda}\left(tb+(1-t)a, \lambda d+(1-\lambda)c\right)\right|^q \\
\leq\ & t^s\lambda^s\left|\frac{\partial^2 f}{\partial t \partial \lambda}\right|^q(b,d)+t^s(1-\lambda)^s\left|\frac{\partial^2 f}{\partial t \partial \lambda}\right|^q(b,c) \\
&+\lambda^s(1-t)^s\left|\frac{\partial^2 f}{\partial t \partial \lambda}\right|^q(a,d)+(1-\lambda)^s(1-t)^s\left|\frac{\partial^2 f}{\partial t \partial \lambda}\right|^q(a,c)
\end{aligned}$$

thus, we obtain

$$\begin{aligned}
&\left|\frac{f(a,c)+r_2 f(a,d)+r_1 f(b,c)+r_1 r_2 f(b,d)}{(r_1+1)(r_2+1)}+\frac{1}{(b-a)(d-c)}\int_a^b\int_c^d f(x,y)dxdy\right. \\
&-\left(\frac{r_2}{r_2+1}\right)\frac{1}{d-c}\int_c^d f(b,y)dy-\left(\frac{1}{r_1+1}\right)\frac{1}{d-c}\int_c^d f(a,y)dy \\
&\left.-\left(\frac{r_2}{r_2+1}\right)\frac{1}{b-a}\int_a^b f(x,d)dx-\left(\frac{1}{r_2+1}\right)\frac{1}{b-a}\int_a^b f(x,c)dx\right| \\
=\ & \frac{(b-a)(d-c)}{(r_1+1)(r_2+1)}\left(\frac{\left(1+r_1^{\frac{p+1}{p}}\right)\left(1+r_2^{\frac{p+1}{p}}\right)}{(r_1+1)^{\frac{1}{p}}(r_2+1)^{\frac{1}{p}}(p+1)^{\frac{2}{p}}}\right) \\
&\times\left(\frac{\left|\frac{\partial^2 f}{\partial t \partial \lambda}\right|^q(a,c)+\left|\frac{\partial^2 f}{\partial t \partial \lambda}\right|^q(a,d)+\left|\frac{\partial^2 f}{\partial t \partial \lambda}\right|^q(b,c)+\left|\frac{\partial^2 f}{\partial t \partial \lambda}\right|^q(b,d)}{(s+1)^2}\right)^{\frac{1}{q}}.
\end{aligned}$$

Which completes the proof of the inequality (3.4). □

Corollary 3. *(1) Under the assumptions of Theorem 12, if we choose* $r_1 = r_2 = 1$ *in (3.4), we have*

$$\begin{aligned}
(3.5)\qquad & \left| \frac{f(a,c)+f(a,d)+f(b,c)+f(b,d)}{4} \right. \\
& -\frac{1}{2}\left[\frac{1}{d-c}\int_c^d [f(b,y)+f(a,y)]\,dy + \frac{1}{b-a}\int_a^b [f(x,d)+f(x,c)]\,dx\right] \\
& \left. +\frac{1}{(b-a)(d-c)}\int_a^b\int_c^d f(x,y)dxdy \right| \\
= & \frac{(b-a)(d-c)}{4(p+1)^{\frac{2}{p}}} \\
& \times\left(\frac{\left|\frac{\partial^2 f}{\partial t\partial\lambda}\right|^q(a,c)+\left|\frac{\partial^2 f}{\partial t\partial\lambda}\right|^q(a,d)+\left|\frac{\partial^2 f}{\partial t\partial\lambda}\right|^q(b,c)+\left|\frac{\partial^2 f}{\partial t\partial\lambda}\right|^q(b,d)}{(s+1)^2}\right)^{\frac{1}{q}}.
\end{aligned}$$

(2) Under the assumptions of Theorem 12, if we choose $r_1 = r_2 = 0$ *in (3.4), we have*

$$\begin{aligned}
& \left| f(a,c) - \frac{1}{d-c}\int_c^d f(a,y)dy - \frac{1}{b-a}\int_a^b f(x,c)dx \right. \\
& \left. +\frac{1}{(b-a)(d-c)}\int_a^b\int_c^d f(x,y)dxdy \right| \\
= & \frac{(b-a)(d-c)}{(p+1)^{\frac{2}{p}}} \\
& \times\left(\frac{\left|\frac{\partial^2 f}{\partial t\partial\lambda}\right|^q(a,c)+\left|\frac{\partial^2 f}{\partial t\partial\lambda}\right|^q(a,d)+\left|\frac{\partial^2 f}{\partial t\partial\lambda}\right|^q(b,c)+\left|\frac{\partial^2 f}{\partial t\partial\lambda}\right|^q(b,d)}{(s+1)^2}\right)^{\frac{1}{q}}.
\end{aligned}$$

Remark 4. *If we choose* $s = 1$ *in (3.5), we obtain an improvement for the inequality (1.3).*

Theorem 12. *Let* $f : \Delta = [a,b]\times[c,d] \subset [0,\infty)^2 \to [0,\infty)$ *be an absolutely continuous function on* Δ. *If* $\left|\frac{\partial^2 f}{\partial t\partial\lambda}\right|^q$ *is* $s-$*convex function on the co-ordinates on*

Δ, for some fixed $s \in (0,1]$ and $q \geq 1$, then one has the inequality:

$$\begin{aligned}
&\left| \frac{f(a,c)+r_2 f(a,d)+r_1 f(b,c)+r_1 r_2 f(b,d)}{(r_1+1)(r_2+1)} + \frac{1}{(b-a)(d-c)} \int_a^b \int_c^d f(x,y)dxdy \right. \\
&-\left(\frac{r_2}{r_2+1}\right)\frac{1}{d-c}\int_c^d f(b,y)dy - \left(\frac{1}{r_1+1}\right)\frac{1}{d-c}\int_c^d f(a,y)dy \\
&\left. -\left(\frac{r_2}{r_2+1}\right)\frac{1}{b-a}\int_a^b f(x,d)dx - \left(\frac{1}{r_2+1}\right)\frac{1}{b-a}\int_a^b f(x,c)dx \right| \\
\leq\ & \frac{(b-a)(d-c)}{(r_1+1)(r_2+1)} \left(\frac{\left(1+r_1^2\right)\left(1+r_2^2\right)}{4(r_1+1)(r_2+1)} \right)^{1-\frac{1}{q}} \\
&\times \left(\frac{MN\left|\frac{\partial^2 f}{\partial t \partial \lambda}\right|^q (a,c) + LN\left|\frac{\partial^2 f}{\partial t \partial \lambda}\right|^q (a,d) + KM\left|\frac{\partial^2 f}{\partial t \partial \lambda}\right|^q (b,c) + KL\left|\frac{\partial^2 f}{\partial t \partial \lambda}\right|^q (b,d)}{(s+1)^2 (s+2)^2} \right)^{\frac{1}{q}}
\end{aligned}$$

for some fixed $r_1, r_2 \in [0,1]$.

Proof. From Lemma 3 and using the well-known Power-mean inequality, we can write

$$\begin{aligned}
&\left| \frac{f(a,c)+r_2 f(a,d)+r_1 f(b,c)+r_1 r_2 f(b,d)}{(r_1+1)(r_2+1)} + \frac{1}{(b-a)(d-c)} \int_a^b \int_c^d f(x,y)dxdy \right. \\
&-\left(\frac{r_2}{r_2+1}\right)\frac{1}{d-c}\int_c^d f(b,y)dy - \left(\frac{1}{r_1+1}\right)\frac{1}{d-c}\int_c^d f(a,y)dy \\
&\left. -\left(\frac{r_2}{r_2+1}\right)\frac{1}{b-a}\int_a^b f(x,d)dx - \left(\frac{1}{r_2+1}\right)\frac{1}{b-a}\int_a^b f(x,c)dx \right| \\
=\ & \frac{(b-a)(d-c)}{(r_1+1)(r_2+1)} \left(\int_0^1 \int_0^1 \left|\left((r_1+1)t-1\right)\left((r_2+1)\lambda-1\right)\right| dtd\lambda \right)^{1-\frac{1}{q}} \\
&\times \left(\int_0^1 \int_0^1 \left|\left((r_1+1)t-1\right)\left((r_2+1)\lambda-1\right)\right| \left|\frac{\partial^2 f}{\partial t \partial \lambda}\left(tb+(1-t)a, \lambda d+(1-\lambda)c\right)\right|^q dtd\lambda \right)^{\frac{1}{q}}.
\end{aligned}$$

Since $\left|\frac{\partial^2 f}{\partial t \partial \lambda}\right|^q$ is $s-$convex function on the co-ordinates on Δ, we can write for $t, \lambda \in [0,1]$

$$\begin{aligned}
&\left|\frac{\partial^2 f}{\partial t \partial \lambda}\left(tb+(1-t)a, \lambda d+(1-\lambda)c\right)\right|^q \\
\leq\ & t^s \left|\frac{\partial^2 f}{\partial t \partial \lambda}\left(b, \lambda d+(1-\lambda)c\right)\right|^q + (1-t)^s \left|\frac{\partial^2 f}{\partial t \partial \lambda}\left(a, \lambda d+(1-\lambda)c\right)\right|^q
\end{aligned}$$

and

$$\begin{aligned}
&\left|\frac{\partial^2 f}{\partial t\partial\lambda}\left(tb+(1-t)a,\lambda d+(1-\lambda)c\right)\right|^q\\
\leq\ & t^s\lambda^s\left|\frac{\partial^2 f}{\partial t\partial\lambda}\right|^q(b,d)+t^s(1-\lambda)^s\left|\frac{\partial^2 f}{\partial t\partial\lambda}\right|^q(b,c)\\
&+\lambda^s(1-t)^s\left|\frac{\partial^2 f}{\partial t\partial\lambda}\right|^q(a,d)+(1-\lambda)^s(1-t)^s\left|\frac{\partial^2 f}{\partial t\partial\lambda}\right|^q(a,c)
\end{aligned}$$

hence, it follows that

$$\begin{aligned}
&\left|\frac{f(a,c)+r_2 f(a,d)+r_1 f(b,c)+r_1 r_2 f(b,d)}{(r_1+1)(r_2+1)}\right.\\
&-\left(\frac{r_2}{r_2+1}\right)\frac{1}{d-c}\int_c^d f(b,y)dy-\left(\frac{1}{r_1+1}\right)\frac{1}{d-c}\int_c^d f(a,y)dy\\
&-\left(\frac{r_2}{r_2+1}\right)\frac{1}{b-a}\int_a^b f(x,d)dx-\left(\frac{1}{r_2+1}\right)\frac{1}{b-a}\int_a^b f(x,c)dx \qquad (3.6)\\
&\left.+\frac{1}{(b-a)(d-c)}\int_a^b\int_c^d f(x,y)dxdy\right|\\
\leq\ & \frac{(b-a)(d-c)}{(r_1+1)(r_2+1)}\left(\frac{\left(1+r_1^2\right)\left(1+r_2^2\right)}{4(r_1+1)(r_2+1)}\right)^{1-\frac{1}{q}}\\
&\times\left(\int_0^1\int_0^1\left|\left((r_1+1)t-1\right)\left((r_2+1)\lambda-1\right)\right|\left\{t^s\lambda^s\left|\frac{\partial^2 f}{\partial t\partial\lambda}\right|^q(b,d)\right.\right.\\
&+t^s(1-\lambda)^s\left|\frac{\partial^2 f}{\partial t\partial\lambda}\right|^q(b,c)+\lambda^s(1-t)^s\left|\frac{\partial^2 f}{\partial t\partial\lambda}\right|^q(a,d)\\
&\left.\left.+(1-\lambda)^s(1-t)^s\left|\frac{\partial^2 f}{\partial t\partial\lambda}\right|^q(a,c)\right\}dtd\lambda\right)^{\frac{1}{q}}
\end{aligned}$$

By a simple computation, one can see that

$$\begin{aligned}
&\left(\int_0^1\int_0^1\left|\left((r_1+1)t-1\right)\left((r_2+1)\lambda-1\right)\right|\left\{t^s\lambda^s\left|\frac{\partial^2 f}{\partial t\partial\lambda}\right|^q(b,d)\right.\right.\\
&+t^s(1-\lambda)^s\left|\frac{\partial^2 f}{\partial t\partial\lambda}\right|^q(b,c)+\lambda^s(1-t)^s\left|\frac{\partial^2 f}{\partial t\partial\lambda}\right|^q(a,d)\\
&\left.\left.+(1-\lambda)^s(1-t)^s\left|\frac{\partial^2 f}{\partial t\partial\lambda}\right|^q(a,c)\right\}dtd\lambda\right)^{\frac{1}{q}}\\
=\ & \left(\frac{MN\left|\frac{\partial^2 f}{\partial t\partial\lambda}\right|^q(a,c)+LN\left|\frac{\partial^2 f}{\partial t\partial\lambda}\right|^q(a,d)+KM\left|\frac{\partial^2 f}{\partial t\partial\lambda}\right|^q(b,c)+KL\left|\frac{\partial^2 f}{\partial t\partial\lambda}\right|^q(b,d)}{(s+1)^2(s+2)^2}\right)^{\frac{1}{q}}
\end{aligned}$$

where K, L, M and N as in Theorem 11. By substituting these in (3.6) and simplifying we obtain the required result. □

Corollary 4. *(1) Under the assumptions of Theorem 13, if we choose* $r_1 = r_2 = 1$*, we have*

$$\begin{aligned}
&\left|\frac{f(a,c)+f(a,d)+f(b,c)+f(b,d)}{4}\right.\\
&-\frac{1}{2}\left[\frac{1}{d-c}\int_c^d [f(b,y)+f(a,y)]\,dy+\frac{1}{b-a}\int_a^b [f(x,d)+f(x,c)]\,dx\right]\\
&\left.+\frac{1}{(b-a)(d-c)}\int_a^b\int_c^d f(x,y)dxdy\right|\\
\leq\ & \frac{(b-a)(d-c)}{4}\left(\frac{1}{4}\right)^{1-\frac{1}{q}}\\
&\times\left(\frac{MN\left|\frac{\partial^2 f}{\partial t\partial\lambda}\right|^q(a,c)+LN\left|\frac{\partial^2 f}{\partial t\partial\lambda}\right|^q(a,d)+KM\left|\frac{\partial^2 f}{\partial t\partial\lambda}\right|^q(b,c)+KL\left|\frac{\partial^2 f}{\partial t\partial\lambda}\right|^q(b,d)}{(s+1)^2(s+2)^2}\right)^{\frac{1}{q}}
\end{aligned}$$

(2) Under the assumptions of Theorem 13, if we choose $r_1 = r_2 = 0$*, we have*

$$\begin{aligned}
&\left|f(a,c)+\frac{1}{(b-a)(d-c)}\int_a^b\int_c^d f(x,y)dxdy\right.\\
&\left.-\frac{1}{d-c}\int_c^d f(a,y)dy-\frac{1}{b-a}\int_a^b f(x,c)dx\right|\\
\leq\ & (b-a)(d-c)\left(\frac{1}{4}\right)^{1-\frac{1}{q}}\\
&\times\left(\frac{MN\left|\frac{\partial^2 f}{\partial t\partial\lambda}\right|^q(a,c)+LN\left|\frac{\partial^2 f}{\partial t\partial\lambda}\right|^q(a,d)+KM\left|\frac{\partial^2 f}{\partial t\partial\lambda}\right|^q(b,c)+KL\left|\frac{\partial^2 f}{\partial t\partial\lambda}\right|^q(b,d)}{(s+1)^2(s+2)^2}\right)^{\frac{1}{q}}
\end{aligned}$$

Remark 5. *Under the assumptions of Theorem 13, if we choose* $r_1 = r_2 = 1$ *and* $s = 1$*, we have an improvement for the inequality (1.4).*

References

- [1] N.S. Barnett and S.S. Dragomir, An Ostrowski type inequality for double integrals and applications for cubature formulae, Soochow J. Math., 27 (1) (2001), 1-10.
- [2] S.S. Dragomir, On Hadamard's inequality for convex functions on the co-ordinates in a rectangle from the plane, Taiwanese Journal of Math., 5, 2001, 775-788.
- [3] B.G. Pachpatte, A new Ostrowski type inequality for double integrals, Soochow J. Math., 32 (2) (2006), 317-322.
- [4] M. Z. Sarıkaya, On the Ostrowski type integral inequality for double integral, http://arxiv.org/abs/1005.0454v1.
- [5] M. Alomari and M. Darus, On the Hadamard's inequality for log −convex functions on the coordinates, Journal of Inequalities and Appl., 2009, article ID 283147.
- [6] H. Hudzik and L. Maligranda, Some remarks on $s-$convex functions, Aequationes Math., 48 (1994), 100-111.
- [7] M.K. Bakula and J. Pečarić, On the Jensen's inequality for convex functions on the coordinates in a rectangle from the plane, Taiwanese Journal of Math., 5, 2006, 1271-1292.
- [8] W. Orlicz, A note on modular spaces , I, Bull. Acad. Polon. Sci. Math. Astronom. Phys., 9 (1961), 157-162.

[9] S.S. Dragomir and S. Fitzpatrick, The Hadamard's inequality for $s-$convex functions in the second sense, Demonstratio Math., 32 (4) (1999), 687-696.
[10] M. Alomari and M. Darus, The Hadamard's inequality for $s-$convex functions of 2-variables, Int. Journal of Math. Analysis, 2(13), 2008, 629-638.
[11] M.E. Özdemir, E. Set, M.Z. Sarıkaya, Some new Hadamard's type inequalities for co-ordinated $m-$convex and $(\alpha, m)-$convex functions, Accepted.
[12] M.Z. Sarıkaya, E. Set, M. Emin Özdemir and S.S. Dragomir, New some Hadamard's type inequalities for co-ordinated convex functions, Accepted.
[13] U.S. Kırmacı, M.K. Bakula, M.E. Özdemir and J. Pečarić, Hadamard-type inequalities for $s-$convex functions, Applied Mathematics and Computation, 193 (2007), 26-35.

♦Ataturk University, K.K. Education Faculty, Department of Mathematics, 25240, Erzurum, Turkey
E-mail address: `emos@atauni.edu.tr`

E-mail address: `hkavurmaci@atauni.edu.tr`

♠Ağri İbrahim Çeçen University, Faculty of Science and Arts, Department of Mathematics, 04100, Ağri, Turkey
E-mail address: `ahmetakdemir@agri.edu.tr`

E-mail address: `merveavci@ymail.com`